

Updated NAIS
(for persons age 18 years or older)

Vaccine	18-26 years	27-59 years	60-64 years	≥ 65 years
Influenza (INF)	1 dose annually or per season			1 dose annually or per season
Pneumococcal: • PCV20; <u>or</u> • PCV13 and/or PPSV23	1 dose of PCV20; <u>or</u> 1 dose of PCV13 and/or 1 or more doses of PPSV23, depending on age			or medical condition(s)
Shingles: Recombinant herpes zoster vaccine (RHZV)	2 doses			
COVID-19	1 initial dose (3 doses for immunocompromised), followed by 1 dose annually			
Tetanus, reduced diphtheria and acellular pertussis (Tdap)	1 dose during each pregnancy			
Human papillomavirus (HPV2)	3 doses (Females)			
Hepatitis B (HepB)	3 doses (4 doses for adults with renal insufficiency)			
Measles, mumps and rubella (MMR)	2 doses			
Varicella (VAR)	2 doses			

	Recommended for persons who meet age requirements
	Recommended for persons with specific medical condition or indication
	Recommended for persons who have not been previously vaccinated, or lack evidence of past infection or immunity

Annex B

Updated NCIS (from birth to age 17 years)

Vaccine	Birth	2 months	4 months	6 months	12 months	15 months	18 months	2-4 years	5-9 years	10-11 years	12-13 years	13-14 years	15-17 years
Bacillus Calmette-Guérin (BCG)	D1												
Hepatitis B (HepB)	D1	D2		D3									
Diphtheria, tetanus and acellular pertussis (paediatric) (DTaP)		D1	D2	D3			B1						
Tetanus, reduced diphtheria and acellular pertussis (Tdap)										B2			
Inactivated poliovirus (IPV)		D1	D2	D3			B1			B2			
Haemophilus influenzae type b (Hib)		D1	D2	D3			B1						
Pneumococcal conjugate (PCV13)			D1	D2	B1								
Pneumococcal polysaccharide (PPSV23)								One or two doses for children and adolescents age 2-17 years with specific medical condition or indication.					
Measles, mumps and rubella (MMR)					D1	D2							
Varicella (VAR)					D1	D2							
Human papillomavirus (HPV2)											D1 (Female s)	D2 (Female s)	
Influenza (INF)				Annual vaccination or per season for <u>all children</u> age 6 months to <5 years (6-59 months).					Annual vaccination or per season for children and adolescents age 5-17 years with specific medical condition or indication.				
COVID-19				Two initial doses (three doses for immunocompromised), followed by annual vaccination for children age 6 months to <5 years with specific medical condition or indication.					One initial dose (three doses for immunocompromised), followed by annual vaccination for children and adolescents age 5-17 years with specific medical condition or indication.				

 Recommended ages and doses for all children

 Recommended for persons with specific medical condition or indication

FOOTNOTES:

- **D1, D2, D3:** Dose 1, dose 2, dose 3
- **B1, B2:** Booster 1, booster 2
- **10-11, 12-13, 13-14 years:** Primary 5, Secondary 1, Secondary 2 (Tdap, IPV, HPV (for females) and MMR (as catch-up) vaccines are provided as part of Health Promotion Board's school-based vaccination programme)
- **HepB:** Doses 2 and 3 are recommended to be given as part of the 6-in-1 vaccine at 2 and 6 months, respectively
- **MMR:** Only the dose 2 is recommended to be given as part of the MMRV vaccine