

What should I know about Xafariv?

- Xafariv thins the blood, which prevents you from getting dangerous blood clots.
- Xafariv must be taken exactly as prescribed by your doctor. To ensure optimal protection from blood clots, never skip a dose.
- You must not stop taking Xafariv without first talking to your doctor as your risk of blood clots may increase.

- Tell your healthcare provider about any other medicines you are currently taking, took recently or intend to start taking, before you start Xafariv.
- Tell your healthcare provider that you are taking Xafariv before any surgery or invasive procedure.

When should I seek advice from my healthcare provider?

- When taking a blood thinner such as Xafariv it is important to be aware of its possible side effects.

- Bleeding is the most common side effect. Do not start taking Xafariv if you know you are at risk of bleeding, without first discussing this with your doctor.
- Tell your healthcare provider straight away if you have any signs or symptoms of bleeding such as the following:
 - Pain
 - Swelling or discomfort
 - Headache, dizziness, or weakness
 - Unusual bruising, nosebleeds, bleeding of gums, and cuts that take a long time to stop bleeding

- Menstrual flow or vaginal bleeding that is heavier than normal
- Blood in your urine, which may be pink or brown and red or black stools
- Coughing up blood, or vomiting blood or material that looks like coffee grounds

How do I take Xafariv?

- To ensure optimal protection, Xafariv
 - 2.5 mg can be taken with or without food
 - 10 mg can be taken with or without food
 - 15 mg must be taken with food
 - 20 mg must be taken with food

Xafariv (rivaroxaban) PATIENT ALERT CARD

Xafariv	2.5 mg	☐
Xafariv	10 mg	☐
Xafariv	15 mg	☐
Xafariv	20 mg	☐

(Tick the prescribed dose)

- Keep this card with you at all times
- Present this card to every physician or dentist prior to treatment

I am under anticoagulation treatment with Xafariv (rivaroxaban)

Name:

Other medicines/conditions:

Address:

Birth date:

Weight:

In case of emergency, please notify:

Doctor's name:

Doctor's phone:

Doctor's stamp:

Please also notify:

Name:

Phone:

Relationship:

Information for healthcare providers:

- INR values should not be used as they are not a dependable measure of the anticoagulant activity of Xafariv.

Local version 1.0

This document has been approved by HSA as of 14 Jul 2026.