

In case of emergency,
Please notify:

Doctor's name :

Doctor's phone :

Please also notify:

Name:

Phone :

Relationship:

I am under anticoagulation treatment
with Xabarem (rivaroxaban).

Name:

Address:

Date of birth:

Weight:

Other medications conditions:



Remedica

- Xabarem 2.5 mg film-coated tablets
- Xabarem 10 mg film-coated tablets
- Xabarem 15 mg film-coated tablets
- Xabarem 20 mg film-coated tablets

*Tick the prescribed dose

PATIENT ALERT CARD

- Keep this card with you at all times
- Present this card to every physician or dentist prior to treatment

Information for health care providers:
INR values should not be used as they are not a dependable measure of the anticoagulant activity of Xabarem

What should I know about Xabarem?

- Xabarem thins the blood, which prevents you from getting dangerous blood clots.
- Xabarem must be taken exactly as prescribed by your doctor. To ensure optimal protection from blood clots, never skip a dose.
- You must not stop taking Xabarem without first talking to your doctor as your risk of blood clots may increase.
- Tell your health care provider about any other medicines you are currently taking, took recently or intend to start taking, before you start Xabarem.
- Tell your health care provider that you are taking Xabarem before any surgery or invasive procedure.

When should I seek advice from my health care provider?

- When taking a blood thinner such as Xabarem it is important to be aware of its possible side effects. Bleeding is the most common side effect.
- Do not start taking Xabarem if you know you are at risk of bleeding, without first discussing this with your doctor.
- Tell your health care provider straight away if you have any signs or symptoms of bleeding such as the following:
- pain
 - swelling or discomfort
 - headache, dizziness or weakness
 - unusual bruising, nosebleeds, bleeding of gums, cuts that take a long time to stop bleeding
 - menstrual flow or vaginal bleeding that is heavier than normal
 - blood in your urine which may be pink or brown, red or black stools
 - coughing up blood, or vomiting blood or material that looks like coffee grounds

How do I take Xabarem?

- To ensure optimal protection, Xabarem - 2.5 mg can be taken with or without food
- 10 mg can be taken with or without food
- 15 mg must be taken with food
- 20 mg must be taken with food

This document has been approved by HSA as of 27-03-2026.