

References

This is the reference list to the ACE Clinical Guideline "Chronic kidney disease – early detection".

1. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl.* 2013;3:1-150.
2. GBD Chronic Kidney Disease Collaboration. Global, regional, and national burden of chronic kidney disease, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2020;395:709–733.
3. Epidemiology & Disease Control Division and Policy, Research & Surveillance Group, Ministry of Health and Health Promotion Board, Singapore. National Population Health Survey 2020. Available from: www.moh.gov.sg/resources-statistics/reports/national-survey-2019-20 [Accessed 5 September 2022].
4. Institute for Health Metrics and Evaluation. Singapore 2020. Available from: www.healthdata.org/singapore [Accessed 5 September 2022].
5. Fraser SD, Blakeman T. Chronic kidney disease: and management in primary care. *Pragmat Obs Res.* 2016;7:21-32.
6. Chronic Kidney Disease (CKD) Management in Primary Care (4th edition). Melbourne: Kidney Health Australia; 2020.
7. Gansevoort RT, Matsushita K, van der Velde M et al. Lower estimated GFR and higher albuminuria are associated with adverse kidney outcomes. A collaborative meta-analysis of general and high-risk population cohorts. *Kidney Int.* 2011;80(1):93-104.
8. Van der Velde M, Matsushita K, Coresh J et al. Lower estimated glomerular filtration rate and higher albuminuria are associated with all-cause and cardiovascular mortality. A collaborative meta-analysis of high-risk population cohorts. *Kidney Int.* 2011;79(12):1341-1352.
9. Matsushita K, van der Velde M, Astor BC et al. Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. *Lancet.* 2010;375(9731):2073-2081.
10. Astor BC, Matsushita K, Gansevoort RT et al. Lower estimated glomerular filtration rate and higher albuminuria are associated with mortality and end-stage renal disease. A collaborative meta-analysis of kidney disease population cohorts. *Kidney Int.* 2011;79(12):1331-1340.
11. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl.* 2012;2:1-138.
12. Makris K, Spanou L. Acute kidney injury: definition, pathophysiology and clinical phenotypes. *Clin Biochem Rev.* 2016;37(2):85-98.
13. Lamb EJ, MacKenzie F, Stevens PE. How should proteinuria be detected and measured? *Ann Clin Biochem.* 2009;46(Pt 3):205-217.
14. Ballantyne FC, Gibbons J, O'Reilly DS. Urine albumin should replace total protein for the assessment of glomerular proteinuria. *Ann Clin Biochem.* 1993;30 (Pt 1):101-103.
15. Newman DJ, Thakkar H, Medcalf EA et al. Use of urine albumin measurement as a replacement for total protein. *Clin Nephrol.* 1995;43(2):104-109.
16. Stevens LA, Coresh J, Greene T et al. Assessing kidney function—measured and estimated glomerular filtration rate. *N Engl J Med.* 2006;354(23):2473-2483.
17. Johnson DW, Jones GR, Mathew TH et al. Chronic kidney disease and measurement of albuminuria or proteinuria: a position statement. *Med J Aust.* 2012;197(4):224-225.
18. National Institute for Health and Care Excellence (NICE). Chronic kidney disease in adults: assessment and management. London: National Institute for Health and Care Excellence (UK); 2014.
19. Simerville JA, Maxted WC, Pahira JJ. Urinalysis: a comprehensive review. *Am Fam Physician.* 2005;71(6):1153-1162.
20. Narva AS, Norton JM, Boulware LE. Educating patients about CKD: the path to self-management and patient-centered care. *Clinical journal of the American Society of Nephrology : CJASN.* 2016;11(4):694-703.
21. Hwang SJ, Tan NC, Yoon S et al. Perceived barriers and facilitators to chronic kidney disease care among patients in Singapore: a qualitative study. *BMJ Open.* 2020;10(10):e041788.
22. Johnson DW, Atai E, Chan M et al. KHA-CARI Guideline: Early chronic kidney disease: Detection, prevention and management. *Nephrology.* 2013;18:340-350.
23. Malaysian Health Technology Assessment Section Ministry of Health Malaysia. Management of chronic kidney disease in adults (second edition). Putrajaya: Malaysian Health Technology Assessment Section (MaHTAS), Medical Development Division, Ministry of Health Malaysia; 2018.